



**Virak Tan, MD**  
**Rosemary Yi, MD**  
**Robyn Martin, RN, BSN, CCM**

33 Bleeker Street  
Suite 103  
Millburn, NJ 07041

620 Essex Street  
Suite 202  
Harrison, NJ 07029

Ph: (973) 947-4700  
eFax: (888) 900-9262  
HandArmSurgery.com

---

## Post-Operative Instructions

The following are general instructions that will be useful after your surgery. Unless otherwise instructed, you should follow these instructions.

**Swelling** – This is a natural and expected occurrence after surgery but can cause a great deal of discomfort, as well as, limit joint range of motion and flexibility.

- 1) Elevation will help reduce swelling. This is often accomplished by placing the affected arm on pillows above the level of your heart.
- 2) Ice can be applied directly over your post-operative dressing or cast. Be careful to keep the dressings dry.
- 3) If your fingers are left free move them as often as you can. They may be swollen, but moving them will actually help reduce the swelling. You may use the other hand to assist with finger motion.
- 4) The dressings should be snug. If the dressings feel too tight you can loosen the ace and re-wrap. This may require help from another person.

### Wound Care

For most surgeries, the dressings will remain in place until the first post-operative visit. This is usually within 10-14 days. For some procedures such as trigger finger and carpal tunnel releases, the dressings can be removed on the 5<sup>th</sup> day after surgery. If the incision is healing without oozing or gapping, you may wash the hand gently and shower without scrubbing the surgical site. The incision should not be submerged in water.

### Medication

Pain medicine is usually given post-operatively. Please take as directed. If the medication prescribed is too strong, take half the dose (cut or break the pill in half) or take extra-strength Tylenol. You should resume all the medications that you normally take, unless otherwise instructed.

### Therapy

Therapy is usually prescribed at the first post-operative visit, if needed. There are some procedures that require therapy to begin within the first day or two after surgery. These include, but are not limited to, tendon repair, Dupuytren's release, contracture release and tenolysis. Early therapy is very important in the success of these cases.

**If you have any question, please contact the office.**